



## TG Therapeutics Announces Issuance of Permanent J-Code for BRIUMVI® (ublituximab-xiyy) from Centers for Medicare and Medicaid Services

April 27, 2023

NEW YORK, April 27, 2023 (GLOBE NEWSWIRE) -- TG Therapeutics, Inc. (NASDAQ: TGTX) today announced that the U.S. Centers for Medicare & Medicaid Services (CMS) has issued a permanent J-Code for BRIUMVI (ublituximab-xiyy), for the treatment of adult patients with relapsing forms of multiple sclerosis (RMS). Under the Healthcare Common Procedure Coding System (HCPCS), the BRIUMVI J-Code (J2329) will become effective July 1, 2023.

Michael S. Weiss, the Company's Chairman and Chief Executive Officer, stated, "The issuance of a permanent J-Code for BRIUMVI is a significant milestone in our early launch phase. Once in effect, the permanent J-Code will provide a streamlined and efficient reimbursement process, which we believe will facilitate additional patient access. We are encouraged by the early launch of BRIUMVI and look forward to sharing additional details on our first quarter earnings call."

J-codes are permanent, product specific, reimbursement codes used by commercial insurance plans and government payers for Medicare Part B drugs like BRIUMVI that are administered by a physician. A permanent J-code simplifies the claims submission and documentation process, which in turn streamlines the billing and reimbursement process. The permanent J-code for BRIUMVI is published online on the CMS website [here](#).

### **ABOUT BRIUMVI (ublituximab-xiyy) 150 mg/6 mL Injection for IV**

BRIUMVI is a novel monoclonal antibody that targets a unique epitope on CD20-expressing B-cells. Targeting CD20 using monoclonal antibodies has proven to be an important therapeutic approach for the management of autoimmune disorders, such as RMS. BRIUMVI is uniquely designed to lack certain sugar molecules normally expressed on the antibody. Removal of these sugar molecules, a process called glycoengineering, allows for efficient B-cell depletion.

BRIUMVI is indicated for the treatment of adults with relapsing forms of multiple sclerosis (RMS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease.

A list of authorized specialty distributors can be found at [www.briumvi.com](http://www.briumvi.com).

### **IMPORTANT SAFETY INFORMATION**

**Contraindications: BRIUMVI is contraindicated in patients with:**

- Active HBV infection
- A history of life-threatening infusion reaction to BRIUMVI

### **WARNINGS AND PRECAUTIONS**

**Infusion Reactions:** BRIUMVI can cause infusion reactions, which can include pyrexia, chills, headache, influenza-like illness, tachycardia, nausea, throat irritation, erythema, and an anaphylactic reaction. In MS clinical trials, the incidence of infusion reactions in BRIUMVI-treated patients who received infusion reaction-limiting premedication prior to each infusion was 48%, with the highest incidence within 24 hours of the first infusion. 0.6% of BRIUMVI-treated patients experienced infusion reactions that were serious, some requiring hospitalization.

Observe treated patients for infusion reactions during the infusion and for at least one hour after the completion of the first two infusions unless infusion reaction and/or hypersensitivity has been observed in association with the current or any prior infusion. Inform patients that infusion reactions can occur up to 24 hours after the infusion. Administer the recommended pre-medication to reduce the frequency and severity of infusion reactions. If life-threatening, stop the infusion immediately, permanently discontinue BRIUMVI, and administer appropriate supportive treatment. Less severe infusion reactions may involve temporarily stopping the infusion, reducing the infusion rate, and/or administering symptomatic treatment.

**Infections:** Serious, life-threatening or fatal, bacterial and viral infections have been reported in BRIUMVI-treated patients. In MS clinical trials, the overall rate of infections in BRIUMVI-treated patients was 56% compared to 54% in teriflunomide-treated patients. The rate of serious infections was 5% compared to 3% respectively. There were 3 infection-related deaths in BRIUMVI-treated patients. The most common infections in BRIUMVI-treated patients included upper respiratory tract infection (45%) and urinary tract infection (10%). Delay BRIUMVI administration in patients with an active infection until the infection is resolved.

Consider the potential for increased immunosuppressive effects when initiating BRIUMVI after immunosuppressive therapy or initiating an immunosuppressive therapy after BRIUMVI.

**Hepatitis B Virus (HBV) Reactivation:** HBV reactivation occurred in an MS patient treated with BRIUMVI in clinical trials. Fulminant hepatitis, hepatic failure, and death caused by HBV reactivation have occurred in patients treated with anti-CD20 antibodies. Perform HBV screening in all patients before initiation of treatment with BRIUMVI. Do not start treatment with BRIUMVI in patients with active HBV confirmed by positive results for HBsAg and anti-HB tests. For patients who are negative for surface antigen [HBsAg] and positive for HB core antibody [HBcAb+] or are carriers of HBV [HBsAg+], consult a liver disease expert before starting and during treatment.

**Progressive Multifocal Leukoencephalopathy (PML):** Although no cases of PML have occurred in BRIUMVI-treated MS patients, JCV infection resulting in PML has been observed in patients treated with other anti-CD20 antibodies and other MS therapies.

If PML is suspected, withhold BRIUMVI and perform an appropriate diagnostic evaluation. Typical symptoms associated with PML are diverse, progress over days to weeks, and include progressive weakness on one side of the body or clumsiness of limbs, disturbance of vision, and changes in thinking, memory, and orientation leading to confusion and personality changes.

MRI findings may be apparent before clinical signs or symptoms; monitoring for signs consistent with PML may be useful. Further investigate suspicious findings to allow for an early diagnosis of PML, if present. Following discontinuation of another MS medication associated with PML, lower PML-related mortality and morbidity have been reported in patients who were initially asymptomatic at diagnosis compared to patients who had characteristic clinical signs and symptoms at diagnosis.

If PML is confirmed, treatment with BRIUMVI should be discontinued.

**Vaccinations:** Administer all immunizations according to immunization guidelines: for live or live-attenuated vaccines at least 4 weeks and, whenever possible at least 2 weeks prior to initiation of BRIUMVI for non-live vaccines. BRIUMVI may interfere with the effectiveness of non-live vaccines. The safety of immunization with live or live-attenuated vaccines during or following administration of BRIUMVI has not been studied. Vaccination with live virus vaccines is not recommended during treatment and until B-cell repletion.

**Vaccination of Infants Born to Mothers Treated with BRIUMVI During Pregnancy:** In infants of mothers exposed to BRIUMVI during pregnancy, assess B-cell counts prior to administration of live or live-attenuated vaccines as measured by CD19<sup>+</sup> B-cells. Depletion of B-cells in these infants may increase the risks from live or live-attenuated vaccines. Inactivated or non-live vaccines may be administered prior to B-cell recovery. Assessment of vaccine immune responses, including consultation with a qualified specialist, should be considered to determine whether a protective immune response was mounted.

**Fetal Risk:** Based on data from animal studies, BRIUMVI may cause fetal harm when administered to a pregnant woman. Transient peripheral B-cell depletion and lymphocytopenia have been reported in infants born to mothers exposed to other anti-CD20 B-cell depleting antibodies during pregnancy. A pregnancy test is recommended in females of reproductive potential prior to each infusion. Advise females of reproductive potential to use effective contraception during BRIUMVI treatment and for 6 months after the last dose.

**Reduction in Immunoglobulins:** As expected with any B-cell depleting therapy, decreased immunoglobulin levels were observed. Decrease in immunoglobulin M (IgM) was reported in 0.6% of BRIUMVI-treated patients compared to none of the patients treated with teriflunomide in RMS clinical trials. Monitor the levels of quantitative serum immunoglobulins during treatment, especially in patients with opportunistic or recurrent infections, and after discontinuation of therapy until B-cell repletion. Consider discontinuing BRIUMVI therapy if a patient with low immunoglobulins develops a serious opportunistic infection or recurrent infections, or if prolonged hypogammaglobulinemia requires treatment with intravenous immunoglobulins.

**Most Common Adverse Reactions:** The most common adverse reactions in RMS trials (incidence of at least 10%) were infusion reactions and upper respiratory tract infections.

Physicians, pharmacists, or other healthcare professionals with questions about BRIUMVI should visit [www.briumvi.com](http://www.briumvi.com).

#### **ABOUT BRIUMVI PATIENT SUPPORT**

BRIUMVI Patient Support is a flexible program designed by TG Therapeutics to support patients through their treatment journey in a way that works best for them. More information about the BRIUMVI Patient Support program can be accessed at [www.briumvipatientsupport.com](http://www.briumvipatientsupport.com).

#### **ABOUT MULTIPLE SCLEROSIS**

Relapsing multiple sclerosis (RMS) is a chronic demyelinating disease of the central nervous system (CNS) and includes people with relapsing-remitting multiple sclerosis (RRMS) and people with secondary progressive multiple sclerosis (SPMS) who continue to experience relapses. RRMS is the most common form of multiple sclerosis (MS) and is characterized by episodes of new or worsening signs or symptoms (relapses) followed by periods of recovery. It is estimated that nearly 1 million people are living with MS in the United States and approximately 85% are initially diagnosed with RRMS.<sup>1,2</sup> The majority of people who are diagnosed with RRMS will eventually transition to SPMS, in which they experience steadily worsening disability over time. Worldwide, more than 2.3 million people have a diagnosis of MS.<sup>1</sup>

#### **ABOUT TG THERAPEUTICS**

TG Therapeutics is a fully integrated, commercial stage, biopharmaceutical company focused on the acquisition, development and commercialization of novel treatments for B-cell diseases. In addition to a research pipeline including several investigational medicines, TG has received approval from the U.S. FDA for BRIUMVI (ublituximab-xiyy), for the treatment of adult patients with relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease. For more information, visit [www.tgtherapeutics.com](http://www.tgtherapeutics.com), and follow us on Twitter [@TGTherapeutics](https://twitter.com/TGTherapeutics) and on [LinkedIn](https://www.linkedin.com/company/tgtherapeutics).

#### **Cautionary Statement**

This press release contains forward-looking statements that involve a number of risks and uncertainties. For those statements, we claim the protection of the safe harbor for forward-looking statements contained in the Private Securities Litigation Reform Act of 1995.

Any forward-looking statements in this press release are based on management's current expectations and beliefs and are subject to a number of risks, uncertainties and important factors that may cause actual events or results to differ materially from those expressed or implied by any forward-looking statements contained in this press release. In addition to the risk factors identified from time to time in our reports filed with the U.S. Securities and Exchange Commission (SEC), factors that could cause our actual results to differ materially include the below.

Such forward looking statements include but are not limited to statements regarding expectations for the progression or success of our commercial launch of BRIUMVI (ublituximab-xiyy) for RMS and statements regarding the impact of the issuance of a permanent J-Code for BRIUMVI.

Additional factors that could cause our actual results to differ materially include the following: the Company's ability to maintain a commercial infrastructure for BRIUMVI, and to successfully market and sell BRIUMVI; the impact of the issuance of a permanent J-Code on streamlining the reimbursement process and or facilitating patient access; the failure to obtain and maintain requisite regulatory approvals, including the risk that the Company fails to satisfy post-approval regulatory requirements; the Company's ability to meet post-approval compliance obligations (on topics

including but not limited to product quality, product distribution and supply chain, pharmacovigilance, and sales and marketing); the Company's reliance on third parties for manufacturing, distribution and supply, and other support functions for our clinical and commercial products, including BRIUMVI, and the ability of the Company and its manufacturers and suppliers to produce and deliver BRIUMVI to meet the market demand for BRIUMVI. Further discussion about these and other risks and uncertainties can be found in our Annual Report on Form 10-K for the fiscal year ended December 31, 2022 and in our other filings with the U.S. Securities and Exchange Commission.

Any forward-looking statements set forth in this press release speak only as of the date of this press release. We do not undertake to update any of these forward-looking statements to reflect events or circumstances that occur after the date hereof. This press release and prior releases are available at [www.tgtherapeutics.com](http://www.tgtherapeutics.com). The information found on our website is not incorporated by reference into this press release and is included for reference purposes only.

CONTACT:

**Investor Relations**

Email: [ir@tgtxinc.com](mailto:ir@tgtxinc.com)

Telephone: 1.877.575.TGTX (8489), Option 4

**Media Relations:**

Email: [media@tgtxinc.com](mailto:media@tgtxinc.com)

Telephone: 1.877.575.TGTX (8489), Option 6

---

1. MS Prevalence. National Multiple Sclerosis Society website. <https://www.nationalmssociety.org/About-the-Society/MS-Prevalence>. Accessed October 26, 2020. 2. Multiple Sclerosis International Federation, 2013 via Datamonitor p. 236.